

# Assay Requisition Form

Date Sent: \_\_\_\_\_ Date and Time Received: \_\_\_\_\_

Requested By: \_\_\_\_\_ Title: \_\_\_\_\_

Organization: \_\_\_\_\_ PO# (if applicable): \_\_\_\_\_

Signature: \_\_\_\_\_

Send Results to: \_\_\_\_\_ Bill to: \_\_\_\_\_  Same as Results

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Special Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

ELISA Assays	Catalog No.	Fee	Quantity	SAMPLE ID: _____
<input type="checkbox"/> IL-1 $\beta$	1002	\$72.00	_____	<b>SAMPLE TYPE:</b> <input type="checkbox"/> Cell Culture Media <input type="checkbox"/> Porcine Serum <input type="checkbox"/> Porcine Plasma (specify below) <input type="checkbox"/> Heparin <input type="checkbox"/> EDTA <input type="checkbox"/> Citrate
<input type="checkbox"/> IL-6	1003	\$72.00	_____	
<input type="checkbox"/> IL-10	1004	\$72.00	_____	
<input type="checkbox"/> TGF $\beta$	1005	\$72.00	_____	
<input type="checkbox"/> TNF $\alpha$	1006	\$72.00	_____	

PCR Assays	Catalog No.	Fee	Quantity	SAMPLE ID: _____
<input type="checkbox"/> $\beta$ -actin	2001	\$60.00	_____	<b>SAMPLE TYPE:</b> <input type="checkbox"/> cDNA <input type="checkbox"/> mRNA Specify Preserved Specimen Type: _____
<input type="checkbox"/> IL-1 $\beta$	2002	\$137.00	_____	
<input type="checkbox"/> IL-6	2003	\$137.00	_____	
<input type="checkbox"/> IL-10	2004	\$137.00	_____	
<input type="checkbox"/> TGF $\beta$	2005	\$137.00	_____	
<input type="checkbox"/> TNF $\alpha$	2006	\$137.00	_____	
<input type="checkbox"/> AdipoR1	2007	\$137.00	_____	
<input type="checkbox"/> AdipoR2	2008	\$137.00	_____	

Endotoxin Assays	Catalog No.	Fee	Quantity	SAMPLE ID: _____
<input type="checkbox"/> Gel Clot (LAL)	3001	\$150.00	_____	<input type="checkbox"/> pH 6-8 <input type="checkbox"/> Other pH: _____

**Specimen Requirements:**

**ELISA assays:** Please submit a minimum of 2 mL of frozen sample. Please contact Xeno Diagnostics for further information on alternative sample sizes

**Endotoxin assay:** Please submit a minimum of 5mL of culture media for testing. Alternatively sufficient material for reconstitution in 5mL of pyrogen free water is required. Note that the presence of  $\beta$ -glucans can cause the apparent endotoxin level to be increased. Also the optimal pH for the assay is pH6-8. Please indicate if sample is buffered outside of this pH range.

**PCR assays:** Please contact Xeno Diagnostics for further details.